

A dynamical model for longitudinal wave functions in light-front holographic QCD

Sophia S. Chabysheva and John R. Hiller

Department of Physics
University of Minnesota-Duluth
Duluth, Minnesota 55812
(Dated: August 1, 2012)

Abstract

We construct a Schrödinger-like equation for the longitudinal wave function of a meson in the valence $q\bar{q}$ sector, based on the 't Hooft model for large- N two-dimensional QCD, and combine this with the usual transverse equation from light-front holographic QCD, to obtain a model for mesons with massive quarks. The computed wave functions are compared with the wave function *ansatz* of Brodsky and De Teramond and used to compute decay constants and parton distribution functions. The basis functions used to solve the longitudinal equation may be useful for more general calculations of meson states in QCD.

PACS numbers: 11.25.Tq, 12.38.Lg, 12.39.Ki, 14.40.-n

I. INTRODUCTION

Light-front holographic QCD [1] provides a Schrödinger-like equation for the transverse wave functions of hadrons, based on a hypothesized duality with AdS_5 . The duality arguments lead to the construction of models for a confining transverse potential. The soft-wall model [2] in particular admits analytic solutions for the spectrum and the transverse wave functions, and, what is more, yields a spectrum [3–7] consistent with linear Regge trajectories. However, because the duality relies on the approximate conformal limit of zero-mass quarks, the transverse models do not include dependence on quark masses. Also, the longitudinal wave function is only specified for zero-mass quarks, through a matching of dual form factor calculations [8].

One escape from these restrictions is an *ansatz* by Brodsky and De Teramond [9] that extends the transverse momentum dependence to include the full invariant mass. For meson states, this extends an exponentiation of $k_\perp^2/x(1-x)$, where $\vec{k}_\perp$ is the transverse momentum and x is the longitudinal momentum fraction, to $k_\perp^2/x(1-x) + m_1^2/x + m_2^2/(1-x)$. To be consistent with the zero-mass limit, we take the quark masses m_1 and m_2 to be current quark masses.

An alternative is to assume separation of variables for the meson wave function and then write down a light-front equation for the longitudinal part that includes quark masses and a model potential. The kinetic term will be just $m_1^2/x + m_2^2/(1-x)$, though here, to be consistent with nonrelativistic quark models, the quark masses are constituent masses. The potential term should be confining and should yield longitudinal wave functions consistent with the dual form-factor analysis, in the zero-current-mass limit.

A potential model that achieves this, and is directly related to QCD, is the 't Hooft model obtained in the large- N limit of two-dimensional QCD [10]. There exist nearly exact analytic solutions for the ground state, which can be improved easily with numerical calculations [11–13]. In fact, a natural basis set for the numerical calculation, which uses Jacobi polynomials [13], could be useful for general calculations of meson valence states. The coupling constant in the model is constrained by the requirement that the longitudinal wave function for the pion be consistent with the dual form-factor analysis when the constituent quark masses have appropriate values.

Here we explore this potential-model approach. The details of the model and its solution are discussed in Sec. II. Sample applications are illustrated in Sec. III. A summary and some additional remarks are included in Sec. IV.

II. THE MODEL

In light-front holographic QCD, the meson valence wave function is written in factorized form

$$\psi_{nLS}(x, \zeta, \varphi) = X(x)\phi_{nLS}(\zeta)\frac{1}{\sqrt{2\pi}}e^{iL\varphi}, \quad (2.1)$$

where n is the transverse radial quantum number, L is the orbital angular momentum, S is the spin, and $\zeta \equiv \sqrt{x(1-x)}b_\perp$ is the transverse variable, with $b_\perp$ the transverse position. The transverse wave function ϕ_{nLS} satisfies a Schrödinger-like equation, which in the soft-

wall model, takes the form [1, 2]

$$\left[-\frac{d^2}{d\zeta^2} - \frac{1-4L^2}{4\zeta^2} + \kappa^2\zeta^2 + 2\kappa^2(L+S-1) \right] \phi_{nLS}(\zeta) = (M^2 - M_{\parallel}^2)\phi_{nLS}, \quad (2.2)$$

with $M_{\parallel}^2$ the longitudinal-transverse separation constant. The analytic solutions are two-dimensional oscillator wave functions

$$\phi_{nLS}(x) = \kappa^{1+L} \sqrt{\frac{2n!}{(n+L)!}} \sqrt{\zeta^{2L+1}} e^{-\kappa^2\zeta^2/s} L_n^L(\kappa^2\zeta^2), \quad (2.3)$$

where the L_n^L are associated Laguerre polynomials. The mass spectrum is

$$M^2 = 4\kappa^2(n+L+S/2) + M_{\parallel}^2. \quad (2.4)$$

In our approach, the longitudinal wave function $X(x)$ satisfies its own Schrödinger-like equation

$$\left[\frac{m_1^2}{x} + \frac{m_2^2}{1-x} \right] X(x) + \frac{g^2}{\pi} \mathcal{P} \int dy \frac{(X(x) - X(y))}{(x-y)^2} - CX(x) = M_{\parallel}^2 X(x), \quad (2.5)$$

with $\mathcal{P}$ indicating the principal value and C a constant. This is the equation obtained in the 't Hooft model for large- N two-dimensional QCD [10]. Because the transverse equation already introduces enough quantum numbers, we consider only the ground state of the longitudinal equation; any additional quantum number associated with longitudinal excitations would represent double counting. Also, since the light-meson spectrum is already represented by the transverse equation, the constant C is used to set $M_{\parallel}$ to zero. The net effect is that the additional longitudinal equation is only for determination of the longitudinal wave function and has nothing to say about the spectrum. To recover nonrelativistic quark models in the appropriate limit, we use constituent masses for m_1 and m_2 .

As discussed by 't Hooft [10] and Bergknoff [11], the ground-state wave function $X(x)$ is well approximated by the form $x^{\beta_1}(1-x)^{\beta_2}$. Their analyses of the endpoint behavior show that β_i should satisfy the transcendental equation

$$\frac{m_i^2\pi}{g^2} - 1 + \pi\beta_i \cot \pi\beta_i. \quad (2.6)$$

From the dual form-factor analysis [8], the light-meson wave function should have this form with $\beta_1 = \beta_2 = 1/2$. If we take the up and down quark masses, m_u and m_d , to be equal, the square-root behavior is obtained if $g^2/\pi = m_u^2$, consistent with $\cot \pi/2 = 0$. This fixes the value of the coupling constant. Although the model could be more flexible if g were flavor dependent, we do not consider this.

The exact solution cannot be obtained analytically. Instead, we solve (2.5) numerically by expanding the wave function X in terms of orthonormal basis functions f_n , chosen to include the analytic approximation explicitly. As discussed by Mo and Perry [13], these basis functions are

$$f_n(x) = N_n x^{\beta_1} (1-x)^{\beta_2} P_n^{(2\beta_2, 2\beta_1)}(2x-1), \quad (2.7)$$

with $P_n^{(2\beta_2, 2\beta_1)}$ the Jacobi polynomial of order n . The normalization factor N_n is given by [14]

$$N_n = \sqrt{(2n+2\beta_1+2\beta_2) \frac{n!\Gamma(n+2\beta_1+2\beta_2+1)}{\Gamma(n+2\beta_1+1)\Gamma(n+2\beta_2+1)}}. \quad (2.8)$$

The solution is then represented as

$$X(x) = \sum_n c_n f_n(x). \quad (2.9)$$

For equal-mass cases, the longitudinal equation obeys an $x \leftrightarrow (1-x)$ symmetry, and only the even- n terms will contribute. In general, we find that only a few terms are needed; the $n=0$ term, for which the Jacobi polynomial is constant and $f_0 \propto x^{\beta_1}(1-x)^{\beta_2}$, represents 90% or more of the probability.

The expansion coefficients c_n are obtained by diagonalizing the longitudinal equation in the f_n basis. The matrix representation is

$$\left(\frac{m_1^2}{m_u^2} A_1 + \frac{m_2^2}{m_u^2} A_2 + B \right) \vec{c} = \xi \vec{c}, \quad (2.10)$$

with $\xi \equiv C/m_u^2$ and

$$(A_1)_{nm} = \int_0^1 \frac{dx}{x} f_n(x) f_m(x), \quad (A_2)_{nm} = \int_0^1 \frac{dx}{1-x} f_n(x) f_m(x), \quad (2.11)$$

$$B_{nm} = \int_0^1 dx \mathcal{P} \int_0^1 dy f_n(x) \frac{f_m(x) - f_m(y)}{(x-y)^2}. \quad (2.12)$$

Following 't Hooft [10], the matrix representation is made explicitly symmetric by rewriting the potential term as

$$B_{nm} = \frac{1}{2} \int_0^1 dx \int_0^1 dy \frac{f_n(x) - f_n(y)}{x-y} \frac{f_m(x) - f_m(y)}{x-y}. \quad (2.13)$$

In addition to the explicit symmetry, which simplifies the matrix diagonalization, this rearrangement also resolves the principal value prescription. The matrices are small because the number of terms needed in the expansion are few; the diagonalization is then straightforward.

III. SAMPLE CALCULATIONS

To see the implications of our model, we compare the form of the longitudinal wave function with the *ansatz* by Brodsky and De Teramond [9], both directly and through the computation of parton distribution functions and decay constants, for the pion, kaon, and J/Ψ . Where parameter values are needed, we use the current-quark parameterization of Vega *et al.* [4] with no additional fits or adjustments. The parameter values are listed in Table I.

For the ground state, the complete wave function is

$$\psi(x, \zeta) = N X(x) e^{-\kappa^2 \zeta^2 / 2} \quad (3.1)$$

or, in terms of the transverse coordinate $b_\perp$,

$$\psi(x, b_\perp) = N X(x) e^{-\kappa^2 x(1-x) b_\perp^2 / 2}. \quad (3.2)$$

TABLE I. Meson parameters and decay constants. All dimensionful parameters are in units of MeV. Results are compared between our model, which uses constituent-quark masses, and the *ansatz* of Brodsky and De Teramond [9], where we use current-quark masses. Parameter and experimental values are from Vega *et al.* [4] and the Particle Data Group [15].

	model		<i>ansatz</i>		decay constant				
meson	m_1	m_2	m_1	m_2	$P_{q\bar{q}}$	κ	model	<i>ansatz</i>	exper.
pion	330	330	4	4	0.204	951	131	132	130
kaon	330	500	4	101	1	524	160	162	156
J/ Ψ	1500	1500	1270	1270	1	894	267	238	278

The factor N is fixed by the normalization

$$P_{q\bar{q}} = \int_0^1 dx \int_0^\infty db_\perp^2 |\psi(x, b_\perp)|^2, \quad (3.3)$$

where $P_{q\bar{q}}$ is the probability of the quark-antiquark valence state. If $X(x)$ is separately normalized such that

$$\int_0^1 dx \frac{|X(x)|^2}{x(1-x)} = 1, \quad (3.4)$$

then $N = \frac{\kappa}{\pi} \sqrt{P_{q\bar{q}}}$. A Fourier transform to transverse momentum $\vec{k}_\perp$ yields

$$\psi(x, k_\perp) = \sqrt{P_{q\bar{q}}} \frac{4\pi}{\kappa} \frac{X(x)}{x(1-x)} e^{-k_\perp^2/(2\kappa^2 x(1-x))}. \quad (3.5)$$

For zero-mass quarks, $X(x) = \sqrt{x(1-x)}$, in order to satisfy the form factor duality [8]. The *ansatz* for $X(x)$ by Brodsky and De Teramond [9] extends this to massive quarks by allowing $k_\perp^2/x(1-x)$ to become the invariant mass $(m_1^2 + k_\perp^2)/x + (m_2^2 + k_\perp^2)/(1-x)$ with the addition of the two mass terms. This gives

$$X_{\text{BdT}}(x) = N_{\text{BdT}} \sqrt{x(1-x)} e^{-(m_1^2/x + m_2^2/(1-x))/2\kappa^2}, \quad (3.6)$$

with

$$N_{\text{BdT}} = \left[\int_0^1 dx e^{-(m_1^2/x + m_2^2/(1-x))/\kappa^2} \right]^{-1/2}. \quad (3.7)$$

Figure 1 compares this *ansatz* with the $X(x)$ computed in our model. For the pion, the two wave functions are essentially the same, since both involve only tiny variations from $\sqrt{x(1-x)}$.

The wave functions can be used to compute decay constants, as given by [16]

$$f_M = 2\sqrt{6} \int_0^1 dx \int_0^\infty \frac{dk_\perp^2}{16\pi^2} \psi(x, k_\perp) \quad (3.8)$$

The results are included in Table I. The values for the pion and kaon are consistent and in agreement with experiment. The value for the J/ Ψ is significantly closer to experiment than the value obtained from the longitudinal *ansatz*.




FIG. 1. Longitudinal wave functions $X(x)$ for the (a) pion, (b) kaon, and (c) J/Ψ . The solid lines are wave functions from our model; the dashed lines show the *ansatz* by Brodsky and De Tera-
mond [9].

We can also compute parton distributions $f(x)$. As discussed in [4] and shown in [17], if the wave function takes the form

$$\psi(x, k_{\perp}) = \eta(x) e^{-k_{\perp}^2 / (2\kappa^2 x(1-x))}, \quad (3.9)$$

the parton distribution is given by

$$f(x) = \frac{\kappa^2}{16\pi^2} x(1-x) \eta^2(x). \quad (3.10)$$

Applying this to our model, we obtain

$$f(x) = P_{q\bar{q}} \frac{X^2(x)}{x(1-x)}. \quad (3.11)$$

For the *ansatz*, we have

$$f_{\text{BdT}}(x) = N_{\text{BdT}}^2 P_{q\bar{q}} e^{-(m_1^2/x + m_2^2/(1-x))/\kappa^2}. \quad (3.12)$$

These are plotted for comparison in Figure 2. The rough similarity of the wave functions translates into similar parton distributions.




FIG. 2. Same as Fig. 1 but for parton distributions $f(x)$ multiplied by x .

IV. CONCLUDING REMARKS

We have constructed and solved a relativistic light-front equation for the longitudinal wave functions of mesons with massive quarks, to be used in tandem with the transverse equation of light-front holographic QCD. Comparisons with the *ansatz* [9] (3.6) show that for lighter mesons, the longitudinal wave functions are quite similar. However, for the J/Ψ ,

there is a notable difference, which translates into a better estimate of the decay constant, as listed in Table I. Wave functions for the J/Ψ , and also the pion and kaon, are shown in Fig. 1, and parton distributions are shown in Fig. 2. The similarity of the longitudinal wave functions provides the *ansatz* with a connection to the fundamental interactions of QCD.

Perhaps the most broadly useful outcome of this exercise is to illustrate that there is a convenient set of basis functions for longitudinal wave functions of meson valence states. These are the f_n defined in (2.7), with the parameters β_i to be optimized as needed for a given application. One could, of course, use the eigenfunctions of the 't Hooft model, but these do not have an analytic form and would add an extra layer of complication to any calculation.

The importance of the choice of basis functions is in the rate of convergence as the basis is expanded. For the alternative of discrete light-cone quantization, it is known that convergence can be much slower [18]. Thus, these basis functions may prove useful for calculations, such as those described in [19], that use the transverse light-front holographic eigenfunctions.

ACKNOWLEDGMENTS

This work was supported in part by the Department of Energy through Contract No. DE-FG02-98ER41087.

-
- [1] S.J. Brodsky and G.F. de Teramond, Phys. Rev. Lett. **96**, 201601 (2006).
 - [2] A. Karch, E. Katz, D.T. Son, and M.A. Stephanov, Phys. Rev. D **74**, 015005 (2006); L. Da Rold and A. Pomarol, Nucl. Phys. B **721**, 79 (2005).
 - [3] S.S. Gershtein, A.K. Likhoded, and A.V. Luchinsky, Phys. Rev. D **74**, 016002 (2006).
 - [4] A. Vega, I. Schmidt, T. Branz, T. Gutsche, and V.E. Lyubovitskij, Phys. Rev. **80**, 055014 (2009).
 - [5] D. Ebert, R.N. Faustov, and V.O. Galkin, Eur. Phys. J. C **66**, 197 (2010).
 - [6] T. Branz, T. Gutsche, V.E. Lyubovitskij, I. Schmidt, and A. Vega, Phys. Rev. D **82**, 074022 (2010); T. Gutsche, V.E. Lyubovitskij, I. Schmidt, and A. Vega, Phys. Rev. D **85**, 076003 (2012).
 - [7] T.M. Kelley, S.P. Bartz, and J. Kapusta, Phys. Rev. D **83**, 016002 (2011).
 - [8] S.J. Brodsky and G.F. de Teramond, Phys. Rev. D **77**, 056007 (2008).
 - [9] S.J. Brodsky and G.F. de Teramond, arXiv:0802.0514.
 - [10] G. 't Hooft, Nucl. Phys. B **75**, 461 (1974).
 - [11] H. Bergknoff, Nucl. Phys. B **122**, 215 (1977).
 - [12] Y. Ma and J.R. Hiller, J. Comput. Phys. **82**, 229 (1989).
 - [13] Y. Mo and R.J. Perry, J. Comput. Phys. **108**, 159 (1993).
 - [14] M. Abramowitz and I.A. Stegun (eds.), *Handbook of Mathematical Functions* (Dover, New York, 1965).
 - [15] K. Nakamura et al.(Particle Data Group), J. Phys. G **37**, 075021 (2010).
 - [16] S.J. Brodsky, T. Huang, and G.P. Lepage, *Proceedings of the Banff Summer Institute on Particles and Fields 2, Banff, Alberta, 1981*, edited by A.Z. Capri and A.N. Kamal (Plenum,

- New York, 1983), P. 143; G.P. Lepage, S.J. Brodsky, T. Huang, and P.B. Mackenzie, *ibid.*, p. 83; T. Huang, AIP Conf. Proc. **68**, 1000 (1980).
- [17] A.V. Radyushkin, Phys. Rev. D **58**, 114008 (1998).
- [18] B. van de Sande, Phys. Rev. D **54**, 6347 (1996).
- [19] J.P. Vary *et al.*, Phys. Rev. C **81**, 035205 (2010).